

The possibility of quasi-bound state formation of η -meson with helium isotopes

V. A. Tryasuchev* and A. V. Isaev
Tomsk Polytechnic University, Tomsk, Russia

The necessary conditions of quasi-bound state formation of η -meson with isotopes ^3He , ^4He have been found within the framework of optical potential model. These conditions have been compared with the findings about helium nucleus densities and with the available information about ηN -scattering length. Thus, we have conclude that within the framework of discussed model η - ^3He quasi-bound state formation is not possible, but η - ^4He quasi-bound state formation is possible with the great probability.

PACS numbers: 21.10.-k

The interaction of η -mesons with helium isotopes has been considered in the frame of optical potential for the purpose of η - ^3He and η - ^4He quasi-bound state formation. Let us connect the optical potential distributions with the nuclear densities of the discussed nuclei using the findings of root-mean-square radii. For the description of low energy ηN -interaction let us use a well-known fact of resonance domination $S_{11}(1535)$ in the amplitude of this interaction at such energies. In that case the optical potential of η - A interaction of $U(r)$ takes the following form [1]:

$$2\mu U(r) = -4\pi(1 + \frac{m_\eta}{m_N})\rho(r)a_0, \quad (1)$$

where, m_η , m_N are the meson and nucleon masses, μ is the reduced meson-nucleus mass, a_0 is the ηN -scattering length, $\rho(r)$ is the spherically symmetrical density of nucleon in nuclei, which has been chosen in Fermi form:

$$\rho(r) = \frac{\rho_0}{1 + \exp(\frac{r-R_c}{a})}. \quad (2)$$

Here R_c is the half-density radius, a is the thickness of nucleus diffusion surface layer, ρ_0 is the nucleon density of nucleus in the center. For the nucleus with the nucleon number A , two parameters in distribution (2) may can be fixed by the conditions:

$$A = \int_0^\infty r^2 \rho(r) dr, \quad \langle r^2 \rangle = \frac{1}{A} \int_0^\infty r^4 \rho(r) dr, \quad (3)$$

where $\langle r^2 \rangle^{1/2}$ is the root-mean-square radius (r_{rms}) of the nucleus. The knowledge of rms radii of ^3He , ^4He nuclei [2] leaves only one free parameter of $U(r)$ radial distribution, which is called “diffuseness” and stand for a/R_c . The nucleus densities depend on the diffuseness parameters, as it may be seen in figures 1, 2.

For the formation of quasi-bound state in the complex potential with complex energy eigenvalue $E = -(\varepsilon + i\frac{\Gamma}{2})$ where ε is the binding energy, and Γ is the level width,

the definite relation between absolute values of imaginary and real parts of this potential is required, at which the bound state is possible [3, 4]. The calculated formation boundaries, $\varepsilon \approx 0$, of the discussed η -nuclei in the dependence of imaginary potential part on the real one for different nucleon distributions in nuclei ^3He and ^4He are shown in figures 3, 4 in the complex plane of free ηN -scattering length. It is evident that at the nucleus diffuseness decrease for the quasi-bound state formation the greater real potential part is required, that is, real parts of ηN -scattering length. At the diffuseness increase, when a/R_c is over 0.25, the formation boundaries of η -nuclei practically stop shifting to the left, that limits the dependence of quasi-bound state formation on the nuclear density distribution. In view of impossibility of complex ηN -scattering length experimental determination a_0 is found indirectly, that is, they are model dependent and differ greatly from paper to paper [5, 6, 7, 8, 9, 10, 11, 12] (see table 1 in paper [13]). The values of ηN -scattering length getting into the darkened areas in figures 3, 4 demonstrate the possibility of η -nucleus formation at the attraction potential initiated by such a_0 . And on the contrary, if these values are left in the white parts of figures 3, 4 the formation is not possible.

The bound states spectrum simulating results in η - ^3He system are contradictory [13, 14, 15] and ambiguous [16]. The same situation presents in calculations of bound states in η - ^4He system [13, 17]. One may see in figure 3 that formation of quasi-bound state η - ^3He for the known data concerning length of ηN -scattering and root-mean-square radius of ^3He nucleus is impossible for any diffuseness of ^3He nucleus density. If we use “off-shell” length of ηN -scattering in potential (1) as authors of paper [13] insist, the conclusion will be the same because the real and imaginary parts of a_0 should, according to the cited paper, decrease proportionally. On the contrary, the existence of quasi-bound state of η - ^4He within the limits of used model is possible and is almost independent from the diffuseness of ^4He nucleus density as it may be seen in figure 4 if $|Re(a_0)| \geq 0.60$ fm.

The experimental result for both reactions $d + d \rightarrow ^4\text{He} + \eta$, $d + p \rightarrow ^3\text{He} + \eta$ near the thresholds point

*Electronic address: trs@npi.tpu.ru




FIG. 1: The nucleon density distributions of ${}^3\text{He}$ nucleus for different diffuseness parameter values a/R_c and fixed root-mean-square radii. The curve parameters are given in table 1.

TABLE I: Curves for nucleus ${}^3\text{He}$

r_{rms} (fm)	N_-^0	a/R_c	R_c (fm)	ρ_0 (nucleon/fm 3)
1.9	1	0.10	2.210	0.060
	2	0.15	1.991	0.074
	3	0.20	1.770	0.093
	4	0.25	1.571	0.114




FIG. 2: The nucleon density distributions of ${}^4\text{He}$ nucleus for different diffuseness parameter values a/R_c and fixed root-mean-square radii. The curve parameters are given in table 2.

TABLE II: Curves for nucleus ${}^4\text{He}$

r_{rms} (fm)	N_-^0	a/R_c	R_c (fm)	ρ_0 (nucleon/fm 3)
1.6	1	0.05	2.021	0.113
	2	0.10	1.874	0.132
	3	0.15	1.687	0.163
	4	0.20	1.499	0.203
	5	0.25	1.331	0.250

the features. The reaction's cross-sections beginning at the thresholds do not increase with increase of final particle's momenta [17, 18]. This fact does not contradict to the formation of quasi-bound states of η -meson with these nuclei, but such features of cross-section dependencies may be explained by the poles in the amplitudes of $\eta^3\text{He}$ - and $\eta^4\text{He}$ -scattering, that are located directly near the reaction's thresholds. Our conclusion concerning the impossibility of formation of bound states in η - ${}^3\text{He}$ system in accord with calculation result of $\eta^3\text{He}$ -interaction in many-particle approach [14] but contradicts to the conclusion of theoretical papers [13, 15] in which the systems of η - ${}^3\text{He}$ in order to find bound states were considered using different approximations. The result of the latest experiment from COZY [19] doesn't approve the possibility of η -nucleus of ${}^3\text{He}$ system formation. Another conclusion concerning a possibility of bound state formation in η - ${}^4\text{He}$ system does not contradict to result of paper [13] but in our case it is possible at larger values of real parts of ηN -scattering length.

In the conclusion we may say that basing on the exact solution of Schrodinger equation with optical potential (1) and Woods-Saxon radial dependence that bound state η - ${}^4\text{He}$ is possible, but one of η - ${}^3\text{He}$ is not. The features of virtual state of η - ${}^3\text{He}$ have been observed in the experiment [18], while it may be that experiment [17] the features of bound state of η - ${}^4\text{He}$ have been observed.

Authors thank for help in the work L. Sukhikh.




FIG. 3: Curves are the boundaries of quasi-bound states in system η - ^3He . The darkened areas are the areas of quasi-bound state formation of η -meson with ^3He nucleus in the complex plane of ηN -scattering length for different diffuseness a/R_c parameters: 1 – 0.25; 2 – 0.15; 3 – 0.1. ηN -scattering lengths have been taken from works: ■ – [5]; ● – [6]; ▲ – [7]; ▽ – [8]; ◆ – [9]; □ – [10]; ▼ – [11]; ○ – [12].




FIG. 4: Curves are the boundaries of quasi-bound states in system η - ^4He . The darkened areas are the areas of quasi-bound state formation of η -meson with ^4He nucleus in the complex plane of ηN -scattering length for different diffuseness a/R_c parameters: 1 – 0.25; 2 – 0.15; 3 – 0.05. ηN -scattering lengths have been taken from works: ■ – [5]; ● – [6]; ▲ – [7]; ▽ – [8]; ◆ – [9]; □ – [10]; ▼ – [11]; ○ – [12].

-
- [1] T. E. O. Ericson and F. Scheck, Nucl. Phys. **B 19**, 450 (1970).
 [2] L. R. B. Elton, *Nuclear sizes*, (Oxford university press, Oxford) (1961).
 [3] V. A. Tryasuchev, Phys. Atom. Nucl. **60**, 187 (1997).

- [4] A. Sibirtsev, et. al, Phys. Rev. **C 70**, 047001 (2004), nucl-th/0407073.
 [5] M. Arima, et. al, Nucl. Phys. **A 543**, 613 (1992).
 [6] A. M. Green and S. Wycech, Phys. Rev. **C 55**, R2167 (1997).

- [7] M. Batinić and A. Švarc, Few-Body Systems **20**, 69 (1996).
- [8] N. Kaiser, et. al, Phys. Lett. **B 362**, 23 (1995).
- [9] V. V. Abaev and B. M. K. Nefkens, Phys. Rev. **C 53**, 385 (1996).
- [10] C. Sauerman, et. al, Phys. Lett. **B 341**, 261 (1995).
- [11] C. Wilkin, Phys. Rev. **C 47**, R938 (1993).
- [12] L. Tiator, et. al, Nucl. Phys. **A 580**, 455 (1994).
- [13] Q. Haider and L. C. Liu, Phys. Rev. **C 66**, 045208 (2002).
- [14] A. Fix and H. Arenhövel, Phys. Rev. **C 68**, 044002 (2003).
- [15] N. G. Kelkar, et. al, J. Phys. **G 32**, L19 (2006), nucl-th/0601080.
- [16] C. Wilkin, et. al, nucl-ex/0707.1489.
- [17] A. Budzanowski, et. al, (GEM Collaboration), nucl-ex/0811.3372.
- [18] T. Mersmann, et. al, Phys. Rev. Lett. **98**, 242301 (2007).
- [19] W. Krzemien, et. al, nucl-ex/0810.2330.